

# Is a Young Person Trying to Tell You Something?



FAQ

# Q: What are some of the common warning signs of mental health or substance abuse problems among young people?

A: Some of the most common warning signs are:

Doesn't care about favorite activities or is "too tired to	Sudden drop in grades
play"	Changes in friends, especially with drug users
Acts angry and hostile toward people and surroundings	Big changes in personality
Fights or argues with others	Seems to worry constantly about personal safety or
Has trouble sitting still or concentrating most days	safety of family members
Runs away from home	Does not want to go to school
Shows significant weight loss or weight gain	Complains often of headaches, stomachaches and other
Indicates a fear that "bad things will happen"	sicknesses
Talks about death/suicide	Has trouble sleeping or nightmares
Gives away prized personal belongings	Has prolonged feelings of sadness and loneliness

However, it is important to remember that only a licensed medical doctor or mental health professional can provide an assessment and evaluation to determine whether or not someone has a mental health condition or substance abuse disorder.

#### Q: What other conditions and disorders are associated with substance abuse disorder among young people?

A: Conduct disorder (persistent misbehavior) and depression are the two most often reported disorders among young people with substance abuse issues. Post-traumatic stress disorder (PTSD), anxiety disorders, and eating disorders are also common. It is known as a "co-occurring disorder" when someone has both a mental health condition and a substance use disorder.

#### Q: Does the mental health condition cause the substance use disorder when a young person has both?

A: Co-occurring disorders can develop when a person with a mental health condition attempts to cope by taking drugs or alcohol; a behavior often referred to as "self-medicating." It is also common for co-occurring disorders to develop in the opposite order, with alcohol or other drug use causing or exacerbating mental health conditions.

#### Q: What are some of the substances commonly used and abused by youth?

A:

- Tobacco and alcohol
- Street drugs and party drugs like marijuana (pot, weed), cocaine, heroin, GHB (Gamma hydroxybutyrate), synthetic cannabinoids (Spice, K2), Ecstasy, methamphetamine, Cheese (a combination of heroin and cold medication) and LSD
- Prescription drugs such as Ritalin, Ketamine hydrochloride, hydrocodone, OxyContin, Vicodin, Percocet, Xanax, Rohypnol (Roofies), and Steroids
- Over-the-counter drugs like cough, cold, sleep, laxatives and diet medicines
- Inhalants like paint thinners, gasoline, acetone, Freon, glues, correction fluids, felt-tip marker fluid, butane lighters and propane tanks, whipped cream aerosols or dispensers, spray paints, hair or deodorant sprays, canned air and fabric protector sprays

#### Q: What kinds of treatment work for mental health conditions and substance abuse disorders?

A: Seeking treatment for substance use or mental health issues can reduce a young person's risk for further problems, including suicide. Effective treatments for substance use disorders and mental health conditions include individual and group therapy, support groups, and medication. It is also helpful to work with a treatment team of people that talk to each other.

The keys to successful treatment of co-occurring mental health and substance use issues are to treat both simultaneously and to develop an individualized treatment plan using the following treatment approaches (see reverse side):



- O Integrated Treatment Design treats the mental health condition and substance use at the same time. Service providers work together to develop a treatment plan and to coordinate all services, including counseling, case management, medications, social skills training, and family intervention.
- O Cognitive Behavioral Therapy looks at how a person's behavior affects how they feel. This is done by teaching new skills and techniques that give them more control over feelings and reactions.
- Multisystemic Therapy is an intensive, family-based treatment that uses family strengths to encourage positive social behavior and change substance abuse behaviors of a child at home, in school and in public.

## When A Young Person May Need Help

The warning signs for mental health or substance abuse problems among young people listed above are also indicative of a higher risk for suicide. Additional suicide warning signs include:

- Threatening to hurt or kill oneself or talking about wanting to hurt or kill oneself
- ▶ Looking for ways to kill oneself by seeking access to firearms, available pills, or other means
- ▶ Talking or writing about death, dying, or suicide when these actions are out of the ordinary for the person
- ▶ Talking about feeling hopeless
- Withdrawing from friends, family, and society
- ▶ Acting reckless or engaging in risky activities

If you suspect that someone you know is thinking about suicide, the best way to approach the situation is to ask.

Asking is the first step in saving a life and is an important way to show that you are hearing them and that you will listen. You can ask the question directly (examples: "Have you thought about suicide?" "Do you want to kill yourself?" "Are you thinking about suicide?"), or indirectly (examples: "Sometimes when people are sad as you are, they think about suicide. Have you ever thought about it?" "Do you ever want to go to bed and never wake up?"). Once you've asked, follow the steps in the blue box.

Find a private area to talk and seek to establish a relationship (if it is someone you don't already know well). Comment on what you see and observe nonjudgmentally and be curious about the perceived problem (how long they've thought about suicide, if they've attempted in the past). Seek to find out if they are at immediate high risk of suicide (see below). Find out to who and where they normally go for help and if they have a regular doctor, mental health provider or counselor (if you don't already know this information). Always be sure to be aware of your own non-verbal reactions and tone of voice.

If you feel that someone is at risk for suicide, take all signs seriously and refer him or her to a mental health professional or one of the additional resources below.

If you feel that someone is at a high risk for suicide (he or she has a plan and/ or access to means such as pills or a gun), do not leave the person alone. Call 911, seek immediate help from a mental health provider at school or in the community, call a mobile crisis outreach team, or take him or her to the nearest hospital emergency room.

### **Mental Health Resources**

National Suicide Prevention Lifeline: www.suicidepreventionlifeline.org • 1-800-273-Talk (8255)

School Counselors, Doctors, Nurses, Mental Health Professionals, Religious Leaders

Mental Health America of Texas www.mhatexas.org

Texas Suicide Prevention Council www.TexasSuicidePrevention.org

Texas Youth Hotline 1-800-210-2278

Boys Town 24/7 Hotline (for all teens, parents, and families) 1-800-448-3000 American Foundation for Suicide Prevention www.afsp.org

Texas Department of State Health Services Community Centers

www.dshs.state.tx.us/mentalhealth.shtm

National Institute of Mental Health www.nimh.nih.gov

Mental Health America www.MentalHealthAmerica.net 1-800-969-NMHA

National Youth Crisis Hotline 1-800-HIT-HOME **Help Lines**Dial 211 or 1-800-833-5948

National Alliance for the Mentally III of Texas 1-800-633-3760 www.namitx.nami.org

American Association of Child and Adolescent Psychiatry www.aacap.org

www.aacap.org 1-800-333-7636

Texas Underage Drinking Hotline 1-888-843-8222